

Probing seesaw at LHC

Borut Bajc¹, Miha Nemevšek¹ and Goran Senjanović²

¹ *J. Stefan Institute, 1001 Ljubljana, Slovenia*

² *International Centre for Theoretical Physics, Trieste, Italy*

ABSTRACT

We have recently proposed a simple SU(5) theory with an adjoint fermionic multiplet on top of the usual minimal spectrum. This leads to the hybrid scenario of both type I and type III seesaw and it predicts the existence of the fermionic SU(2) triplet between 100 GeV and 1 TeV for a conventional GUT scale of about 10^{16} GeV, with main decays into W (Z) and leptons, correlated through Dirac Yukawa couplings, and lifetimes shorter than about 10^{-12} sec. These decays are lepton number violating and they offer an exciting signature of $\Delta L = 2$ dilepton events together with 4 jets at future pp ($p\bar{p}$) colliders. Increasing the triplet mass endangers the proton stability and so the seesaw mechanism could be directly testable at LHC.

1 Introduction

The seesaw mechanism [1] has been recognized as the most natural scenario for understanding the smallness of neutrino mass. It implies the existence of heavy particles, which after being integrated out, lead to the gauge invariant operator [2]

$$\mathcal{L}_{eff} = y_{eff} \frac{LLHH}{M} , \quad (1)$$

with $M \gg M_W$ usually assumed. As shown in [3], there are three different types of heavy particles that can induce (1):

I) SM fermionic singlets, coupled to leptons through Dirac Yukawa couplings and usually called right-handed neutrinos (type I seesaw) [1];

II) SU(2) bosonic triplet ($Y = 2$) coupled to leptons through Majorana type couplings (type II seesaw) [4];

III) SU(2) fermionic triplet ($Y = 0$) coupled to leptons through Dirac Yukawas, just like the singlet ones in I) (type III seesaw) [5].

Whatever one chooses, one needs a predictive theory above the SM in order to shed some light on neutrino masses; otherwise, one can as well stick to the effective operator in (1). The best bet for such a theory is grand unification since it can predict new mass scale(s). It turns out that both type I and type II seesaw find their natural role in SO(10) theory due to the automatically present left-right symmetry [6, 7, 8, 9]. Although SO(10) is sufficient by itself to determine all the parameters in the I) and II) cases, and even the 1-3 mixing angle [10], the check of the seesaw is only indirect: one can at best relate neutrino properties to proton decay. The main point is that both right-handed neutrinos and the SU(2) scalar triplet are predicted to be very heavy, close to the GUT scale.

What about the type III seesaw? It is clearly custom fit for the SU(5) theory, as suggested recently [11], since it only requires adding the adjoint fermions 24_F to the existing minimal model with three generations of quarks and leptons, and 24_H and 5_H Higgs fields. This automatically leads to the hybrid scenario of both type I and type III seesaw, since 24_F has also a SM singlet fermion, i.e. the right-handed neutrino. One ends up with a realistic spectrum of two massive and one massless light neutrino. The massless one can of course pick up a tiny mass due to say Planck scale effects [12] or running effects [13], too small to play any direct phenomenological role.

The main prediction of this theory is the lightness of the fermionic triplet (for a recent alternative scenario with light triplets see [14]). For a conventional value of $M_{GUT} \approx 10^{16}$ GeV, the unification constraints strongly suggest its mass below TeV, relevant for the future colliders such as LHC. The triplet fermion decay predominantly into W (or Z) and leptons, with lifetimes shorter than about 10^{-12} sec.

Equally important, the decays of the triplet are dictated by the same Yukawa couplings that lead to neutrino masses and thus one has an example of predicted low-energy seesaw directly testable at colliders and likely already at LHC.

In this expanded version of the original work we systematically study the spectrum and the couplings of the theory. In the next section we focus on the unification constraints on the particle spectrum. We perform a numerical study using two-loop RGE taking into account various mass scales of the theory. We discuss $b - \tau$ unification and the predictions of the fermionic triplet mass depending on the GUT scale. We find a maximal value of the GUT scale: $M_{GUT} \leq 10^{16}$ GeV, which offers a great hope of observing proton decay in a not so distant future. The color octets turn out not to be light

enough for direct observation.

In section III we focus on the phenomenological implications of the theory for LHC. We discuss carefully the decay modes of the triplets and their connection with neutrino masses and mixings. Whereas for generic values of Yukawa couplings it is not easy to make clear predictions, for the case of vanishing θ_{13} or large Yukawa couplings (possibly related to large flavour violating processes) one can constraint the relevant branching ratios and thus directly test the seesaw mechanism at colliders.

Next, in section IV we turn our attention to cosmology and discuss leptogenesis. We find that it can work only in the resonant regime which implies the same mass of the fermionic triplet and singlet and further constrains the parameters space of the theory. The nice feature of a high degree of predictivity of this theory has also a negative implication: we show that there is no stable particle candidate for the dark matter of the universe. We conclude our work with section V, where we also discuss the relevance of our work for supersymmetry.

2 Unification constraints and the mass scales of the theory

The minimal implementation of the type III seesaw in nonsupersymmetric SU(5) requires a fermionic adjoint 24_F in addition to the usual field content 24_H , 5_H and three generations of fermionic 10_F and $\bar{5}_F$. The consistency of the charged fermion masses requires higher dimensional operators in the usual Yukawa sector [15]. One must add the new Yukawa interactions

$$\begin{aligned} \mathcal{L}_{Y\nu} &= y_0^i \bar{5}_F^i 24_F 5_H \\ &+ \frac{1}{\Lambda} \bar{5}_F^i \left(y_1^i 24_F 24_H + y_2^i 24_H 24_F + y_3^i \text{Tr} 24_F 24_H \right) 5_H + h.c. . \end{aligned} \quad (2)$$

After the SU(5) breaking

$$\langle 24_H \rangle = \frac{M_{GUT}}{\sqrt{30}} \text{diag} (2, 2, 2, -3, -3) \quad (3)$$

one obtains the following physical relevant Yukawa interactions for neutrino with the triplet $\sigma_3^F \equiv \vec{\sigma}_3^F \vec{\tau}$ (type III) and singlet σ_0^F (type I) fermions:

$$\mathcal{L}_{Y\nu} = L_i \left(y_T^i \sigma_3^F + y_S^i \sigma_0^F \right) H + h.c. , \quad (4)$$

where y_T^i, y_S^i are two different linear combinations of y_0^i and $y_a^i M_{GUT}/\Lambda$ ($a = 1, 2, 3$). It is clear from the above formula that besides the new appearance of the triplet fermion, the singlet fermion in 24_F acts precisely as the right-handed neutrino; it should not come out as a surprise, as it has the right SM quantum numbers.

Even before we discuss the physical consequences in detail, one important prediction emerges: only two light neutrinos get mass, while the third one remains massless.

In order to discuss the masses of the new fermions, we need the new Yukawa couplings between 24_F and 24_H

$$\begin{aligned} \mathcal{L}_F &= m_F \text{Tr} 24_F^2 + \lambda_F \text{Tr} 24_F^2 24_H \\ &+ \frac{1}{\Lambda} \left(a_1 \text{Tr} 24_F^2 \text{Tr} 24_H^2 + a_2 (\text{Tr} 24_F 24_H)^2 \right. \\ &\left. + a_3 \text{Tr} 24_F^2 24_H^2 + a_4 \text{Tr} 24_F 24_H 24_F 24_H \right) , \end{aligned} \quad (5)$$

where we include the higher dimensional terms for the sake of consistency. The masses of the new fermions are

$$m_0^F = m_F - \frac{\lambda_F M_{GUT}}{\sqrt{30}} + \frac{M_{GUT}^2}{\Lambda} \left[a_1 + a_2 + \frac{7}{30} (a_3 + a_4) \right] , \quad (6)$$

$$m_3^F = m_F - \frac{3\lambda_F M_{GUT}}{\sqrt{30}} + \frac{M_{GUT}^2}{\Lambda} \left[a_1 + \frac{3}{10} (a_3 + a_4) \right] , \quad (7)$$

$$m_8^F = m_F + \frac{2\lambda_F M_{GUT}}{\sqrt{30}} + \frac{M_{GUT}^2}{\Lambda} \left[a_1 + \frac{2}{15} (a_3 + a_4) \right] , \quad (8)$$

$$m_{(3,2)}^F = m_F - \frac{\lambda_F M_{GUT}}{2\sqrt{30}} + \frac{M_{GUT}^2}{\Lambda} \left[a_1 + \frac{(13a_3 - 12a_4)}{60} \right] . \quad (9)$$

Next we turn to the bosonic sector of the theory. We will need the potential for the heavy field 24_H

$$V_{24_H} = m_{24}^2 \text{Tr} 24_H^2 + \mu_{24} \text{Tr} 24_H^3 + \lambda_{24}^{(1)} \text{Tr} 24_H^4 + \lambda_{24}^{(2)} \left(\text{Tr} 24_H^2 \right)^2 , \quad (10)$$

and its interaction with the light fields

$$\begin{aligned}
V_{5_H} = & m_H^2 5_H^\dagger 5_H + \lambda_H \left(5_H^\dagger 5_H \right)^2 + \mu_H 5_H^\dagger 24_H 5_H \\
& + \alpha 5_H^\dagger 5_H Tr 24_H^2 + \beta 5_H^\dagger 24_H^2 5_H .
\end{aligned} \tag{11}$$

It is a straightforward exercise to show that the masses of the bosonic triplet and octet are arbitrary and that one can perform the doublet-triplet splitting through the usual fine-tuning. However splitting its mass from the triplet and the octet fermion masses require the inclusion of higher dimensional terms, which in turn gives an upper bound to the mass of the leptoquark

$$m_{(3,2)}^F \lesssim \frac{M_{GUT}^2}{\Lambda} , \tag{12}$$

where Λ is the cutoff of the theory. One could take naively Λ on the order of the Planck scale, since the theory is asymptotically free. However, without higher dimensional operators one predicts $m_b = m_\tau$ at the GUT scale [16], which fails badly, as much as in the standard model, and thus one must take a lower cut-off¹. To see this we did a one-loop Yukawa running, with a two-loop gauge running. The result $y_\tau \approx 0.01$ and the ratio $y_b/y_\tau \lesssim 0.65$ is valid for any physically allowed value of M_{GUT} .

Thus the analysis requires a cut-off at most two orders of magnitude above the GUT scale. In what follows we take $\Lambda = 100 M_{GUT}$ to ensure the correct mass relations and maximize perturbativity (for a lower cutoff see [17]).

We are now fully armed to study the constraints on the particle spectrum by performing the renormalization group analysis. For the sake of illustration we present first the one-loop analysis. From [11] one has

$$\exp \left[30\pi \left(\alpha_1^{-1} - \alpha_2^{-1} \right) (M_Z) \right] = \tag{13}$$

$$\begin{aligned}
& \left(\frac{M_{GUT}}{M_Z} \right)^{84} \left(\frac{(m_3^F)^4 m_3^B}{M_Z^5} \right)^5 \left(\frac{M_{GUT}}{m_{(3,2)}^F} \right)^{20} \left(\frac{M_{GUT}}{m_T} \right) , \\
& \exp \left[20\pi \left(\alpha_1^{-1} - \alpha_3^{-1} \right) (M_Z) \right] =
\end{aligned} \tag{14}$$

¹We thank Ilja Doršner for pointing it to us. For further details see [17].

$$\left(\frac{M_{GUT}}{M_Z}\right)^{86} \left(\frac{(m_8^F)^4 m_8^B}{M_Z^5}\right)^5 \left(\frac{M_{GUT}}{m_{(3,2)}^F}\right)^{20} \left(\frac{M_{GUT}}{m_T}\right)^{-1},$$

where $m_3^{F,B}$, $m_8^{F,B}$, $m_{(3,2)}^F$ and m_T are the masses of weak triplets, colour octets, (only fermionic) leptoquarks and (only bosonic) colour triplets respectively.

From the well known problem in the standard model of the low meeting scale of α_1 and α_2 , it is clear that the SU(2) triplet should be as light as possible and the colour triplet as heavy as possible. In order to illustrate the point, take $m_3^F = m_3^B = M_Z$ and $m_T = M_{GUT}$. This gives $(\alpha_1^{-1}(M_Z) = 59, \alpha_2^{-1}(M_Z) = 29.57, \alpha_3^{-1}(M_Z) = 8.55) M_{GUT} \approx 10^{15.5}$ GeV. Increasing the triplet masses $m_3^{F,B}$ reduces M_{GUT} dangerously, making proton decay too fast.

For more reliable results one needs a two-loop analysis. We focused on the following regions in parameter space:

- 1) $m_8^{F,B} > 10^5$ GeV to comply with cosmological bounds coming from nucleosynthesis. This limit is analogous to the limit on the sfermion masses in split supersymmetry [18, 19] coming from gluino lifetime [20]. At the time of nucleosynthesis all colour octets should have already decayed into a righthanded quark and an off-shell colour triplet through the Yukawa interactions (2);
- 2) $m_T > 10^{12}$ GeV from proton decay;
- 3) $M_{GUT} > 10^{15.5}$ GeV again from proton decay;
- 4) $m_{(3,2)}^F < M_{GUT}^2/\Lambda = M_{GUT}/100$ from (12) and the above discussion on the choice of the cutoff.

The two-loop analysis maintain an approximate dependence on the combinations $m_3 \equiv ((m_3^F)^4 m_3^B)^{1/5}$ and $m_8 \equiv ((m_8^F)^4 m_8^B)^{1/5}$ as at 1-loop order (13), (14). This is useful in the numerical analysis, since one can first use as varying parameters just these combinations, and the extrapolate the result for the case of different fermionic and bosonic masses.

We have seen that at 1-loop order the mass of the fermionic triplet is predicted to lie below TeV. This bound gets somewhat relaxed at 2-loop order, as can be seen from Fig. 1.

The fermionic triplet can be even higher at the price of lowering the bosonic triplet. It must be stressed although, that these maximal values are not typical: one must stretch the parameters, i.e. go to some corner in




Figure 1: The maximum value of the effective triplet mass m_3 as a function of the unification scale M_{GUT} from the two-loop analysis.

parameter space to evade the 1-loop bounds. In other words, in most of the parameter space the bound $m_3^F \lesssim \text{TeV}$ still persists.

It has been noticed in [17], that the constraint 4) for $m_{(3,2)}^F$ can actually be evaded. In fact, there are solutions, in which $m_{(3,2)}^F \approx m_8^F/2$ that can be of order M_{GUT} . We have been however unable to find any solution with M_{GUT} bigger than 10^{15} GeV, which makes them less realistic due to likely problems in large proton decay widths.

Finally, one can ask, where must the octets be. Taking $M_{GUT} = 10^{15.5}$ GeV one can find the possible region in $m_3 - m_8$ plane, that leads to unification (different solutions for m_8 for the same m_3 correspond to different choices of $m_{(3,2)}^F$). This region is shown in Fig. 2.

3 Phenomenological implications: testing seesaw at LHC?

In the previous section we learned that the triplets are quite light, even likely to be found at LHC. How would they be identified?

The Yukawa couplings of the triplet and singlet fermion are parametrized by (we choose the basis in which the Dirac Yukawa matrix between e^c and L is diagonal and real, while y_T^i are real)




Figure 2: The region that gives unification at $M_{GUT} = 10^{15.5}$ GeV.

$$\begin{aligned}
\mathcal{L}_Y &= -y_E^i H^\dagger e_i^c L_i + y_T^i H^T i\tau^2 \tau^a T^a L_i + y_S^i H^T i\tau^2 S L_i + h.c. \\
&= -\frac{v+h}{\sqrt{2}} \left[y_E^i e_i^c e_i + y_T^i \left(\sqrt{2} T^+ e_i + T^0 \nu_i \right) + y_S^i S \nu_i \right] + h.c. \quad (15)
\end{aligned}$$

where $T^\pm$, T^0 are the three states from the fermionic triplet, while S is the fermionic singlet. We have changed the cumbersome notation from the previous section (where it was necessary), since this whole section is devoted only to the fermionic triplet and singlet.

The Majorana masses for the triplet and singlet (with properly defined T^k and S the masses m_T and m_S can be made real and positive) are

$$\mathcal{L}_m = -\frac{m_T}{2} (2T^+ T^- + T^0 T^0) - \frac{m_S}{2} S S + h.c. \quad (16)$$

To the leading order in the neutrino Dirac Yukawa couplings the following transformations define the physical states:

$$\nu_j \rightarrow \nu_j + \epsilon_T^j T^0 + \epsilon_S^j S, \quad (17)$$

$$T^0 \rightarrow T^0 - \epsilon_T^k \nu_k, \quad (18)$$

$$S \rightarrow S - \epsilon_S^k \nu_k, \quad (19)$$

$$e_j \rightarrow e_j + \sqrt{2} \epsilon_T^j T^-, \quad (20)$$

$$T^- \rightarrow T^- - \sqrt{2} \epsilon_T^k e_k, \quad (21)$$

$$T^+ \rightarrow T^+, \quad e^c \rightarrow e^c \quad (22)$$

where

$$\epsilon_X^i \equiv \frac{y_X^i v}{\sqrt{2} m_X} . \quad (23)$$

In the above equation recall that T^+ is a different state from T^- , just like e^c is a different state from e .

The light neutrino mass matrix is then given by

$$m_{ij}^\nu = -\frac{v^2}{2} \left(\frac{y_T^i y_T^j}{m_T} + \frac{y_S^i y_S^j}{m_S} \right) \quad (24)$$

in the basis in which the charged Yukawas and the couplings with W are diagonal.

3.1 $T \rightarrow W(Z) + \text{light lepton}$

These are the predominant decay modes of the triplets, whose strength is dictated by the neutral Dirac Yukawa couplings.

$$\Gamma(T^- \rightarrow Z e_k^-) = \frac{m_T}{32\pi} |y_T^k|^2 \left(1 - \frac{m_Z^2}{m_T^2}\right)^2 \left(1 + 2\frac{m_Z^2}{m_T^2}\right) , \quad (25)$$

$$\sum_k \Gamma(T^- \rightarrow W^- \nu_k) = \frac{m_T}{16\pi} \left(\sum_k |y_T^k|^2\right) \left(1 - \frac{m_W^2}{m_T^2}\right)^2 \left(1 + 2\frac{m_W^2}{m_T^2}\right) , \quad (26)$$

$$\begin{aligned} \Gamma(T^0 \rightarrow W^+ e_k^-) &= \Gamma(T^0 \rightarrow W^- e_k^+) = \\ &= \frac{m_T}{32\pi} |y_T^k|^2 \left(1 - \frac{m_W^2}{m_T^2}\right)^2 \left(1 + 2\frac{m_W^2}{m_T^2}\right) , \end{aligned} \quad (27)$$

$$\sum_k \Gamma(T^0 \rightarrow Z \nu_k) = \frac{m_T}{32\pi} \left(\sum_k |y_T^k|^2\right) \left(1 - \frac{m_Z^2}{m_T^2}\right)^2 \left(1 + 2\frac{m_Z^2}{m_T^2}\right) , \quad (28)$$

where we averaged over initial polarizations and summed over final ones. From (27) one sees that the decays of T^0 , just as those of righthanded neutrinos, violate lepton number. In a machine such as LHC one would typically produce a pair $T^+ T^0$ (or $T^- T^0$), whose decays then allow for interesting $\Delta L = 2$ signatures of same sign dileptons and 4 jets. This fairly SM background free signature is characteristic of any theory with righthanded neutrinos as discussed in [21]. The main point here is that these triplets are

really predicted to be light, unlike in the case of righthanded neutrinos. The detailed analyses of the LHC signatures including the production, the decays and the background is now in progress [22].

The decay rates above are rather sensitive to the Yukawa couplings which on the other hand can vary a lot. First of all, they are not directly related to the neutrino properties, and they are of course rather flavour dependent. The dominant rate goes through the largest Yukawa coupling which has an approximate lower limit of $\approx 5 \times 10^{-7}$ from the atmospheric neutrino oscillations. This translates into the following upper limit for the lifetime of the dominant two-body triplet decay, for say $m_T \approx 300$ GeV

$$\tau_T \lesssim 10^{-1} \text{ mm} . \quad (29)$$

Measuring the above decays means in some sense checking the seesaw parameters. Let's see in more detail this correspondence. The situation with the singlet and triplet making the light neutrino massive through the seesaw mechanism is analogous to the situation with two righthanded neutrinos (for a recent review of this situation see [23]). Thus we can use the known relations [24] (in the case of hierarchical case, i.e. $m_1^\nu = 0$)

$$\frac{vy_T^{i*}}{\sqrt{2}} = i\sqrt{m_T} \left(U_{i2}\sqrt{m_2^\nu} \cos z \pm U_{i3}\sqrt{m_3^\nu} \sin z \right) , \quad (30)$$

$$\frac{vy_S^{i*}}{\sqrt{2}} = -i\sqrt{m_S} \left(U_{i2}\sqrt{m_2^\nu} \sin z \mp U_{i3}\sqrt{m_3^\nu} \cos z \right) , \quad (31)$$

or (in the case of inverse hierarchy, i.e. $m_3^\nu = 0$)

$$\frac{vy_T^{i*}}{\sqrt{2}} = i\sqrt{m_T} \left(U_{i1}\sqrt{m_1^\nu} \cos z \pm U_{i2}\sqrt{m_2^\nu} \sin z \right) , \quad (32)$$

$$\frac{vy_S^{i*}}{\sqrt{2}} = -i\sqrt{m_S} \left(U_{i1}\sqrt{m_1^\nu} \sin z \mp U_{i2}\sqrt{m_2^\nu} \cos z \right) , \quad (33)$$

valid in a different basis than used before, since here y_T^i are not necessarily real. To compare with the previous results one needs just to compute the absolute value $|y_T^i|$ in (30), (32).

In the formulae above z is a complex number, while U the PMNS matrix, that diagonalizes the neutrino mass matrix (24) (for the experimental values and limits see [25])

$$m^\nu = U^* \begin{pmatrix} m_1^\nu & 0 & 0 \\ 0 & m_2^\nu & 0 \\ 0 & 0 & m_3^\nu \end{pmatrix} U^\dagger \quad (34)$$

Suppose we could measure from T decays the Yukawa couplings y_T^i . Then, in the above formulae we have the following unknowns: one complex number z and two CP phases, assuming that the 1 – 3 mixing will be measured soon (keep in mind that there is one CP phase less in the case of one massless neutrino). In general it is not possible to give much constraints or to make some nontrivial checks, since one has 3 real measurements (the absolute values of y_T^i), but 4 parameters available to describe them. In some special cases however the above relations simplify and some nontrivial constraints appear.

As an example consider the inverse hierarchical case with a vanishing θ_{13} . One gets

$$\frac{\Gamma(\tau)}{\Gamma(\mu)} = \tan^2 \theta_{atm} \quad (35)$$

independent on the phases. This can serve as a direct test of the theory if the inverse hierarchy and a small enough θ_{13} are to be established in the future.

Another interesting case is $|Im(z)| \gg 1$, which is equivalent to the large Dirac Yukawa limit. Here the complex parameter z disappears from the branching ratios, which then depend only on the in principle measurable parameters of the PMNS mixing matrix.

3.2 $T^\pm \rightarrow T^0$ decays

For a nonvanishing and positive mass split $\Delta m_T \equiv m_{T^+} - m_{T^0}$ the charged triplet fermion can decay into a neutral one and an (off-shell) W .

One gets for Δm_T at the one-loop level

$$\Delta m_T = \frac{\alpha_2}{2\pi} \frac{m_W^2}{m_T} \left[f\left(\frac{m_T}{m_Z}\right) - f\left(\frac{m_T}{m_W}\right) \right], \quad (36)$$

where

$$f(y) = \frac{1}{4y^2} \log y^2 - \left(1 + \frac{1}{2y^2}\right) \sqrt{4y^2 - 1} \arctan \sqrt{4y^2 - 1}, \quad (37)$$

which gives $\Delta m_T \approx 160$ MeV with 10% accuracy in the whole range $m_Z \leq m_T \leq \infty$. Notice that there is also a possible direct tree-level contribution from (5) through a non-vanishing vev of the bosonic triplet

$$\Delta m_T^{tree} \approx y \langle T_B \rangle . \quad (38)$$

However, $\langle T_B \rangle \lesssim 1$ GeV for the W and Z masses and $y \lesssim 10^{-2}$ since suppressed by M_{GUT}/Λ , so $\Delta m_T^{tree} \lesssim 10$ MeV, a negligible addition.

The fastest decay mode through the above mass difference is clearly $T^\pm \rightarrow T^0 \pi^\pm$, estimated to be $\mathcal{O}(10^{-10})$ sec [26], negligible in comparison with the $W^\pm \nu$ or $Z l^\pm$ decay channels considered in the previous subsection.

In short, the triplet mass difference can be safely ignored.

3.3 $T \rightarrow H +$ light lepton

If the fermionic triplet is heavier than the SM Higgs, it can decay unsuppressed also to the Higgs and a light lepton. The decay widths can be calculated from (15) to give

$$\Gamma(T^- \rightarrow h e_k^-) = \frac{m_T}{32\pi} |y_T^k|^2 \left(1 - \frac{m_h^2}{m_T^2}\right)^2 , \quad (39)$$

$$\sum_k \Gamma(T^0 \rightarrow h \nu_k) = \frac{m_T}{32\pi} \left(\sum_k |y_T^k|^2\right) \left(1 - \frac{m_h^2}{m_T^2}\right)^2 . \quad (40)$$

These results can now explain the apparent “puzzle” from the results (25)-(28). In fact, these decays come out to be nonzero also in the SU(2) preserving limit ($v \rightarrow 0$). However, in this limit there is no mixing between the triplets and the light leptons, so apparently no decays. The results (39)-(40) explains the discrepancy: in this limit there are four degrees of freedom from the Higgs doublet and the final states in (25)-(28) should be interpreted as Z being the imaginary partner of the standard Higgs, and W being the complex partner in the doublet (the upper component). It is easy to check that the exact SU(2) gauge symmetry connects the results (25)-(28) with (39)-(40) in the limiting case $v, m_h \rightarrow 0$.

4 Cosmological implications

4.1 Dark matter

As usual, in order to have a viable dark matter candidate, it must be stable for at least the age of the universe, which can be translated into an extremely small decay width:

$$\Gamma \lesssim 10^{-42} \text{ GeV} . \quad (41)$$

Let us systematically consider various possible candidates:

- 1) the fermionic neutral triplet T^0 : obviously this cannot work, see (29);
- 2) next consider the bosonic triplet from 24_H , with a mass of at least $m_T^B \geq 100 \text{ GeV}$ from collider constraints. Now, the following operator

$$\bar{5}_F \frac{24_H}{\Lambda} 10_F 5_H^\dagger \quad (42)$$

is needed to correct the $b-\tau$ unification, as discussed in the previous section. So the bosonic triplet can decay into a fermion antifermion pair with a decay width of

$$\Gamma \approx \left(\frac{m_W}{\Lambda} \right)^2 m_T^B \quad (43)$$

much too fast ($\approx 10^{-32} \text{ GeV}$) even in the unrealistic case of $\Lambda = M_{Pl}$.

3) what about the bosonic singlet in 24_H ? This singlet is nothing else than the field that breaks $SU(5)$, so the validity of the whole approach requires its mass to be larger than the electroweak scale. Then everything of the previous case applies also here, and thus the same negative conclusion.

4) finally the fermionic singlet S in 24_F . At first sight, this could not work for the same reason as for the fermionic triplet T . However, here there are no such tight constraints on its mass from colliders, so in principle it could weigh around keV. In this case its decay rate gets suppressed strongly by the W propagator, giving

$$\Gamma \approx \left(y_S^i G_F \right)^2 m_S^5 , \quad (44)$$

which is slow enough in the keV region as soon as $y_S^i \lesssim 0.1$, needed anyway for unitarity. Unfortunately, in order to have the right amount of dark matter, such a small mass cannot give a sizeable contribution to the neutrino masses (see [27] for details).

4.2 Leptogenesis

This issue has been discussed at length in a fairly complete paper devoted to the phenomenology and cosmology of the type I seesaw mechanism with only two righthanded neutrinos [23]. The assumption (two ν_R) in that case is a prediction of our model (T and S). The bottom line now is that, as opposed to the generic situation with three righthanded neutrinos², there is a true physical lower limit on the scale of leptogenesis of the order 10^{10} GeV. This can be seen from a straightforward derivation of the following expression for the maximal CP asymmetry (assuming a hierarchy $m_T \ll m_S$)

$$\epsilon_{MAX} \approx \frac{1}{16\pi} \frac{m_T (m_3^\nu - m_2^\nu)}{v^2} . \quad (45)$$

It is evident that for $m_T \approx \text{TeV}$ the CP asymmetry is hopelessly small. The only possibility for leptogenesis in this theory is the resonant [31, 32, 33] one. It is not difficult to show that one can get realistic value for the baryon asymmetry. Following [29] the triplet asymmetry (and similarly the singlet one) can be rewritten as

$$\epsilon_T = -\frac{Im(\vec{y}_T \vec{y}_S^*)^2}{2|\vec{y}_T|^2 |\vec{y}_S|^2} \times \frac{2(m_S^2 - m_T^2) m_T \Gamma_S}{(m_S^2 - m_T^2)^2 + m_T^2 \Gamma_S^2} . \quad (46)$$

where Γ_S is the total decay width of the singlet fermion. The last term in the above product can take its maximal value 1 for properly chosen singlet mass, i.e. for $m_S^2 = m_T^2 + m_T \Gamma_S$. The first term can be rewritten using (30)-(33) and depends on the value of the unknown complex parameter z and the parameters (masses, mixing angles and phases) of the light neutrino sector. This term can be numerically even of order one, showing that a very large asymmetry can be obtained. It has to be stressed however that successful resonant leptogenesis does not allow all values of z . For example, if $Re(z)$ or $Im(z)$ is zero, the final result vanishes. It is interesting that the asymmetry generically decreases with large $Im(z)$ values, restricting the allowed region. A more detailed description with the calculation of the efficiency factor and the inclusion of flavour effects [34, 35] is beyond the scope of this paper and is in progress [36]. Preliminary estimates seem to show that these restrictions are more restrictive than the ones coming from flavour changing neutral processes.

²for a way out of the well known Davidson-Ibarra limit [28], see [29, 30]

The requirement of successful leptogenesis thus puts various constraints on the yet unknown model parameters. Although the degeneracy between the singlet and the triplet mass is not of direct meaning for LHC searches (it may be relevant however for processes under study [36] like $\mu \rightarrow e\gamma$), because the singlets will not be easily produced, the constraints on the Yukawas could be tested measuring the different branching ratios in triplet decays.

5 Summary and outlook

We have recently [11] constructed what can be considered a possible minimal realistic SU(5) grand unified theory. Instead of changing the Higgs sector as conventionally done, we have simply added to the minimal model an adjoint fermion representation which gives a hybrid seesaw scenario, a mixture of type I and type III. The fact that the theory is in accord with experiment may not be so surprising; after all one has enlarged its particle sector. What is remarkable is a prediction of a light fermionic SU(2) triplet, with a mass below TeV, and for a large portion of the parameter space in the LHC reach of below 500 GeV. One has a badly needed predictive theory of seesaw mechanism that can be tested at the collider energies.

Whereas it is always possible to imagine a low energy seesaw, predictive theories such as GUTs normally prefer large seesaw scale, close to M_{GUT} . Even when you assume this scale to be low as often done in the type I case, the production of righthanded neutrinos is suppressed by the small (compared to the gauge couplings) Yukawa couplings (for a recent work see [37]). The type II could of course be tested for a low scale, but again that is not what comes out. In a way, type III seesaw, up to now almost not studied at all, may provide a unique possibility of seesaw tested at LHC.

In this longer version of our letter we have carefully studied some phenomenological and cosmological issues in this theory. We have performed a complete two-loop analysis of unification constraints which confirms the lightness of the triplets, and at the same time predicts $M_{GUT} < 10^{16}$ GeV, implying the proton lifetime below 10^{36} years, possibly observable in the next generation of proton decay experiments. We have discussed the decay of the triplets into the charged leptons and neutrinos and shown how they probe directly neutrino Dirac Yukawa couplings. These Dirac Yukawas could be quite large since small neutrino masses can involve cancellations and in this case lead to possibly observable lepton flavor violating processes. This is

a rather interesting topic and deserves a careful investigation in a separate note. We also confirmed here an expected result that only resonant leptogenesis can work due to the low mass of the fermionic triplet. This would also make a fermionic singlet light; but as in the case of pure type I seesaw it is not of direct phenomenological interest. Finally, we also showed that the theory lacks a dark matter candidate. Of course, one could always add a singlet particle and account for the dark matter if necessary, since for a given mass one can choose an appropriate coupling.

It is interesting to compare this model to the supersymmetric $SU(5)$ theory, or the supersymmetric extension of the standard model. After all, the weak triplet fermions correspond to winos, while the color octet fermions in 24_H correspond to gluinos. Now, it is well known that the sfermions do not enter into the renormalization group constraints, at least not at the one-loop level; they can be as heavy as we wish. This, split supersymmetry program: light winos, gluinos and higgsinos still allows for the unification of gauge couplings, as much as in the case of low energy supersymmetry [38, 39, 40, 41]. Our work shows that the situation can be more complex if one is willing to split supersymmetry: one can have higgsinos completely decoupled, and the gluinos in the intermediate region. But then, by interpolation, there is clearly a continuum of solutions with higgsino anywhere from the weak to the Planck scale, and gluinos from the weak to some intermediate scale. The work on this is in progress and will be reported elsewhere [42]. It is interesting though that LHC may see only winos and nothing else if supersymmetry is to be split, similar as in the theory discussed here. The important difference in our case is the fact that the fermionic triplet, an analogue of winos, is directly related to neutrino masses and mixings. It should be viewed as a possible alternative to low energy supersymmetry; instead of not so well defined principle of naturalness, it has direct physical and phenomenological motivation.

Acknowledgements

The work of G.S. was supported in part by the European Commission under the RTN contract MRTN-CT-2004-503369; the work of B.B. and M.N. was supported by the Slovenian Research Agency. B.B. and M.N. thank ICTP for hospitality during the course of this work. We thank Paolo Creminelli, Ilja Doršner, Tsedenbaljir Enkhbat, Alejandra Melfo, Fabrizio Nesti

and Francesco Vissani for discussion, and Marco Cirelli, Michele Frigerio and Alessandro Strumia for pointing out a sign error in eq. (36) of the first version of the manuscript.

References

- [1] P. Minkowski, Phys. Lett. B **67** (1977) 421; T. Yanagida, proceedings of the *Workshop on Unified Theories and Baryon Number in the Universe*, Tsukuba, 1979, eds. A. Sawada, A. Sugamoto; S. Glashow, in *Cargese 1979, Proceedings, Quarks and Leptons* (1979) ; M. Gell-Mann, P. Ramond, R. Slansky, proceedings of the *Supergravity Stony Brook Workshop*, New York, 1979, eds. P. Van Nieuwenhuizen, D. Freeman R. Mohapatra, G. Senjanović, Phys.Rev.Lett. **44** (1980) 912.
- [2] S. Weinberg, Phys. Rev. Lett. **43** (1979) 1566.
- [3] E. Ma, Phys. Rev. Lett. **81** (1998) 1171 [arXiv:hep-ph/9805219].
- [4] M. Magg and C. Wetterich, Phys. Lett. B **94** (1980) 61; G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181** (1981) 287; R. N. Mohapatra and G. Senjanović, Phys. Rev. D **23** (1981) 165.
- [5] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C **44** (1989) 441;
- [6] J. C. Pati and A. Salam, Phys. Rev. D **10** (1974) 275.
- [7] R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11** (1975) 2558.
- [8] G. Senjanović and R. N. Mohapatra, Phys. Rev. D **12** (1975) 1502.
- [9] G. Senjanović, Nucl. Phys. B **153** (1979) 334.
- [10] For a review of the seesaw in the context of SO(10), see e.g. G. Senjanović, arXiv:hep-ph/0501244.
- [11] B. Bajc and G. Senjanović, arXiv:hep-ph/0612029.
- [12] E. K. Akhmedov, Z. G. Berezhiani and G. Senjanović, Phys. Rev. Lett. **69** (1992) 3013 [arXiv:hep-ph/9205230].

- [13] S. Davidson, G. Isidori and A. Strumia, Phys. Lett. B **646** (2007) 100 [arXiv:hep-ph/0611389].
- [14] S. B. Gudnason, T. A. Rytlov and F. Sannino, arXiv:hep-ph/0612230.
- [15] J. R. Ellis and M. K. Gaillard, Phys. Lett. B **88** (1979) 315.
- [16] M. S. Chanowitz, J. R. Ellis and M. K. Gaillard, Nucl. Phys. B **128** (1977) 506.
- [17] I. Doršner and P. Fileviez Perez, arXiv:hep-ph/0612216.
- [18] N. Arkani-Hamed and S. Dimopoulos, JHEP **0506** (2005) 073 [arXiv:hep-th/0405159].
- [19] G. F. Giudice and A. Romanino, Nucl. Phys. B **699** (2004) 65 [Erratum-ibid. B **706** (2005) 65] [arXiv:hep-ph/0406088].
- [20] A. Arvanitaki, C. Davis, P. W. Graham, A. Pierce and J. G. Wacker, Phys. Rev. D **72** (2005) 075011 [arXiv:hep-ph/0504210].
- [21] W. Y. Keung and G. Senjanović, Phys. Rev. Lett. **50** (1983) 1427. For a recent work see for example F. del Aguila, J. A. Aguilar-Saavedra and R. Pittau, J. Phys. Conf. Ser. **53** (2006) 506 [arXiv:hep-ph/0606198].
- [22] A. Arhrib, B. Bajc, I. Puljak and G. Senjanović, work in progress.
- [23] W. l. Guo, Z. z. Xing and S. Zhou, Int. J. Mod. Phys. E **16** (2007) 1 [arXiv:hep-ph/0612033].
- [24] A. Ibarra and G. G. Ross, Phys. Lett. B **591** (2004) 285 [arXiv:hep-ph/0312138].
- [25] A. Strumia and F. Vissani, arXiv:hep-ph/0606054.
- [26] M. Ibe, T. Moroi and T. T. Yanagida, Phys. Lett. B **644** (2007) 355 [arXiv:hep-ph/0610277].
- [27] T. Asaka, S. Blanchet and M. Shaposhnikov, Phys. Lett. B **631** (2005) 151 [arXiv:hep-ph/0503065].
- [28] S. Davidson and A. Ibarra, Phys. Lett. B **535** (2002) 25 [arXiv:hep-ph/0202239].

- [29] T. Hambye, Y. Lin, A. Notari, M. Papucci and A. Strumia, Nucl. Phys. B **695** (2004) 169 [arXiv:hep-ph/0312203].
- [30] M. Raidal, A. Strumia and K. Turzynski, Phys. Lett. B **609** (2005) 351 [Erratum-ibid. B **632** (2006) 752] [arXiv:hep-ph/0408015].
- [31] M. Flanz, E. A. Paschos, U. Sarkar and J. Weiss, Phys. Lett. B **389** (1996) 693 [arXiv:hep-ph/9607310].
- [32] A. Pilaftsis, Phys. Rev. D **56** (1997) 5431 [arXiv:hep-ph/9707235].
- [33] A. Pilaftsis and T. E. J. Underwood, Nucl. Phys. B **692** (2004) 303 [arXiv:hep-ph/0309342].
- [34] S. Pascoli, S. T. Petcov and A. Riotto, Phys. Rev. D **75** (2007) 083511 [arXiv:hep-ph/0609125].
- [35] S. Pascoli, S. T. Petcov and A. Riotto, Nucl. Phys. B **774** (2007) 1 [arXiv:hep-ph/0611338].
- [36] B. Bajc, M. Nemevšek and G. Senjanović, work in progress.
- [37] J. Kersten and A. Y. Smirnov, arXiv:0705.3221 [hep-ph].
- [38] S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. D **24** (1981) 1681.
- [39] L. E. Ibanez and G. G. Ross, Phys. Lett. B **105** (1981) 439.
- [40] M. B. Einhorn and D. R. T. Jones, Nucl. Phys. B **196** (1982) 475.
- [41] W. J. Marciano and G. Senjanović, Phys. Rev. D **25** (1982) 3092.
- [42] B. Bajc, A. Melfo, F. Nesti and G. Senjanović, work in progress.